



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

From the last theorem we see that a regular group cannot be primitive unless it is generated by a single cycle involving a prime number of letters. Since such a group must be primitive we have the following

THEOREM : *The regular primitive groups and the prime numbers have a 1,1 correspondence ; i. e. for each prime number there is one regular primitive group and for each regular primitive group there is one prime number.*

[To be Continued.]

THE CENTROID OF AREAS AND VOLUMES.

By G. B. M. ZERE, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas.

[Continued from February Number.]

II. VOLUMES. Let the density vary as $x^{h-1}y^{k-1}z^{l-1}$. Then

$$\bar{x} = \frac{\iiint x^h y^{k-1} z^{l-1} dx dy dz}{\iiint x^{h-1} y^{k-1} z^{l-1} dx dy dz}, \quad \bar{y} = \frac{\iiint x^{h-1} y^k z^{l-1} dx dy dz}{\iiint x^{h-1} y^{k-1} z^{l-1} dx dy dz},$$

$$\bar{z} = \frac{\iiint x^{h-1} y^{k-1} z^l dx dy dz}{\iiint x^{h-1} y^{k-1} z^{l-1} dx dy dz}.$$

$$\therefore \bar{x} = \frac{a^{h+1} b^k c^l}{(2m+1)(2n+1)(2p+1)} \frac{\Gamma\left\{\frac{h+1}{2}(2m+1)\right\} \Gamma\left\{\frac{k}{2}(2n+1)\right\} \Gamma\left\{\frac{l}{2}(2p+1)\right\}}{\Gamma\left\{\frac{h+1}{2}(2m+1) + \frac{k}{2}(2n+1) + \frac{l}{2}(2p+1) + 1\right\}}$$

$$\frac{a^h b^k c^l}{(2m+1)(2n+1)(2p+1)} \frac{\Gamma\left\{\frac{h}{2}(2m+1)\right\} \Gamma\left\{\frac{k}{2}(2n+1)\right\} \Gamma\left\{\frac{l}{2}(2p+1)\right\}}{\Gamma\left\{\frac{h}{2}(2m+1) + \frac{k}{2}(2n+1) + \frac{l}{2}(2p+1) + 1\right\}}$$

$$\therefore \bar{x} = \frac{\Gamma(hm+m+\frac{h+1}{2})\Gamma(hm+kn+lp+\frac{h+k+l}{2}+1)}{\Gamma(hm+\frac{h}{2})\Gamma(hm+kn+lp+m+\frac{h+k+l+1}{2}+1)} a \dots\dots\dots (C).$$

$$\bar{y} = \frac{\Gamma(kn+n+\frac{k+1}{2})\Gamma(hm+kn+lp+\frac{h+k+l}{2}+1)}{\Gamma(kn+\frac{k}{2})\Gamma(hm+kn+lp+n+\frac{h+k+l+1}{2}+1)} b \dots\dots\dots (D).$$

$$\bar{z} = \frac{\Gamma(lp+p+\frac{l+1}{2})\Gamma(hm+kn+lp+\frac{h+k+l}{2}+1)}{\Gamma(lp+\frac{l}{2})\Gamma(hm+kn+lp+p+\frac{h+k+l+1}{2}+1)} c \dots\dots\dots (E).$$

This gives the centroid of the eighth part of the volume whatever may be the values of h, k, l, m, n, p .

Let $m=n=p$, and also let the density vary as xyz so that $h=k=l=2$.

$$\therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(3m+\frac{3}{2})\Gamma(6m+4)}{\Gamma(2m+1)\Gamma(7m+\frac{3}{2})}.$$

$$\text{Let } m=0, \therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(\frac{3}{2})\Gamma(4)}{\Gamma(1)\Gamma(\frac{3}{2})} = \frac{16}{35}, \text{ for } \left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 + \left(\frac{z}{c}\right)^2 = 1.$$

$$\text{Let } m=1, \therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(\frac{9}{2})\Gamma(10)}{\Gamma(3)\Gamma(\frac{9}{2})} = \frac{2^{13}}{11.13.17.19},$$

$$\text{for } \left(\frac{x}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{b}\right)^{\frac{2}{3}} + \left(\frac{z}{c}\right)^{\frac{2}{3}} = 1.$$

$$\text{Let } m=\frac{3}{2}, \therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(6)\Gamma(13)}{\Gamma(4)\Gamma(15)} = \frac{10}{91},$$

$$\text{for } \left(\frac{x}{a}\right)^{\frac{1}{2}} + \left(\frac{y}{b}\right)^{\frac{1}{2}} + \left(\frac{z}{c}\right)^{\frac{1}{2}} = 1,$$

the centroid of the volume bounded by the positive portion of the co-ordinate planes.

Let $m=n=p$, and let the density be the same throughout the solid so that, $h=k=l=1$

$$\therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(2m+1)\Gamma(3m+\frac{5}{2})}{\Gamma(m+\frac{1}{2})\Gamma(4m+3)}.$$

Let $m=0$, $\therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(1)\Gamma(\frac{5}{2})}{\Gamma(\frac{1}{2})\Gamma(3)} = \frac{3}{8}$, for $\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 + \left(\frac{z}{c}\right)^2 = 1$.

Let $m=1$, $\therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(3)\Gamma(\frac{7}{2})}{\Gamma(\frac{3}{2})\Gamma(7)} = \frac{21}{128}$, for $\left(\frac{x}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{b}\right)^{\frac{2}{3}} + \left(\frac{z}{c}\right)^{\frac{2}{3}} = 1$.

Let $m=\frac{3}{2}$, $\therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(4)\Gamma(7)}{\Gamma(2)\Gamma(9)} = \frac{3}{28}$, for $\left(\frac{x}{a}\right)^{\frac{1}{2}} + \left(\frac{y}{b}\right)^{\frac{1}{2}} + \left(\frac{z}{c}\right)^{\frac{1}{2}} = 1$.

Let $m=n=p$, and let the density vary as xy , so that $h=k=2$, $l=1$,

$$\therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\Gamma(3m+\frac{3}{2})\Gamma(5m+\frac{3}{2})}{\Gamma(2m+1)\Gamma(6m+4)}, \quad \frac{\bar{z}}{c} = \frac{\Gamma(2m+1)\Gamma(5m+\frac{1}{2})}{\Gamma(m+\frac{1}{2})\Gamma(6m+4)}.$$

Let $m=0$, $\therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\Gamma(\frac{3}{2})\Gamma(\frac{3}{2})}{\Gamma(1)\Gamma(4)} = \frac{5\pi}{32}$,

$$\frac{\bar{z}}{c} = \frac{\Gamma(1)\Gamma(\frac{1}{2})}{\Gamma(\frac{1}{2})\Gamma(4)} = \frac{5}{16}, \text{ for } \left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 + \left(\frac{z}{c}\right)^2 = 1.$$

Let $m=1$, $\therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\Gamma(\frac{9}{2})\Gamma(\frac{7}{2})}{\Gamma(3)\Gamma(10)} = \frac{5 \cdot 7 \cdot 11 \cdot 13 \cdot 15\pi}{2^{20}}$,

$$\frac{\bar{z}}{c} = \frac{\Gamma(3)\Gamma(\frac{5}{2})}{\Gamma(\frac{3}{2})\Gamma(10)} = \frac{5 \cdot 11 \cdot 13}{2^{13}}, \text{ for } \left(\frac{x}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{b}\right)^{\frac{2}{3}} + \left(\frac{z}{c}\right)^{\frac{2}{3}} = 1.$$

Let $m=\frac{3}{2}$, $\therefore \frac{\bar{x}}{a} = \frac{\bar{y}}{b} = \frac{\Gamma(6)\Gamma(11)}{\Gamma(4)\Gamma(13)} = \frac{5}{33}$,

$$\frac{\bar{z}}{c} = \frac{\Gamma(4)\Gamma(11)}{\Gamma(2)\Gamma(13)} = \frac{1}{22}, \text{ for } \left(\frac{x}{a}\right)^{\frac{1}{2}} + \left(\frac{y}{b}\right)^{\frac{1}{2}} + \left(\frac{z}{c}\right)^{\frac{1}{2}} = 1.$$

Let $m=n=p$, and let the density vary as x so that $h=2$, $k=l=1$.

$$\therefore \frac{\bar{x}}{a} = \frac{\Gamma(3m + \frac{3}{2})\Gamma(4m + 3)}{\Gamma(2m + 1)\Gamma(5m + \frac{7}{2})}, \quad \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(2m + 1)\Gamma(4m + 3)}{\Gamma(m + \frac{1}{2})\Gamma(5m + \frac{7}{2})}.$$

Let $m=0$, then for $\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 + \left(\frac{z}{c}\right)^2 = 1$

$$\frac{\bar{x}}{a} = \frac{\Gamma(\frac{3}{2})\Gamma(3)}{\Gamma(1)\Gamma(\frac{7}{2})} = \frac{8}{15}, \quad \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(1)\Gamma(3)}{\Gamma(\frac{1}{2})\Gamma(\frac{7}{2})} = \frac{16}{15\pi}.$$

Let $m=1$, then for $\left(\frac{x}{a}\right)^{\frac{8}{3}} + \left(\frac{y}{b}\right)^{\frac{8}{3}} + \left(\frac{z}{c}\right)^{\frac{8}{3}} = 1$

$$\frac{\bar{x}}{a} = \frac{\Gamma(\frac{8}{3})\Gamma(7)}{\Gamma(3)\Gamma(\frac{17}{3})} = \frac{2^7}{3 \cdot 11 \cdot 13}, \quad \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(3)\Gamma(7)}{\Gamma(\frac{5}{3})\Gamma(\frac{17}{3})} = \frac{2^{14}}{5 \cdot 7 \cdot 9 \cdot 11 \cdot 13\pi}.$$

Let $m=\frac{3}{2}$, then for $\left(\frac{x}{a}\right)^{\frac{1}{2}} + \left(\frac{y}{b}\right)^{\frac{1}{2}} + \left(\frac{z}{c}\right)^{\frac{1}{2}} = 1$

$$\frac{\bar{x}}{a} = \frac{\Gamma(6)\Gamma(9)}{\Gamma(4)\Gamma(11)} = \frac{2}{9}, \quad \frac{\bar{y}}{b} = \frac{\bar{z}}{c} = \frac{\Gamma(4)\Gamma(9)}{\Gamma(2)\Gamma(11)} = \frac{1}{15}.$$

Thus we could multiply examples almost without number.

If $a=b$ we get another series of areas.

If $a=b=c$ we get another series of solids.

If $b=c$ or $a=c$ we get still another series of solids.

But formulæ (A), (B), (C), (D), (E) apply to them all.

One more example and we will proceed to the discussion of surfaces. Let the density vary as x^3y^2z , and let the equation to the surface be

$$\left(\frac{x}{a}\right)^{\frac{8}{3}} + \left(\frac{y}{b}\right)^{\frac{8}{3}} + \left(\frac{z}{c}\right)^{\frac{8}{3}} = 1,$$

so that $h=4$, $k=3$, $l=2$, $m=1$, $n=2$, $p=3$

$$\therefore \bar{x} = \frac{\Gamma(\frac{15}{2})\Gamma(\frac{43}{2})}{\Gamma(6)\Gamma(23)} a = \frac{5 \cdot 7 \cdot 9 \cdot 11 \cdot 13 \cdot 23 \cdot 29 \cdot 31 \cdot 37 \cdot 39 \cdot 41 \pi a}{2^{50}}$$

$$\bar{y} = \frac{\Gamma(10)\Gamma(\frac{43}{2})}{\Gamma(\frac{15}{2})\Gamma(24)} b = \frac{5 \cdot 9 \cdot 29 \cdot 31 \cdot 37 \cdot 41 b}{11 \cdot 2^{26}},$$

$$\bar{z} = \frac{\Gamma(\frac{21}{2})\Gamma(\frac{43}{2})}{\Gamma(7)\Gamma(25)} c = \frac{5 \cdot 7 \cdot 9 \cdot 11 \cdot 13 \cdot 13 \cdot 17 \cdot 19 \cdot 29 \cdot 31 \cdot 37 \cdot 41 \pi c}{2^{51}}.$$

The prodigious amount of work to accomplish this by the ordinary method would be impossible.

[To be Continued.]